

- 3 陈小英,陈知本.金黄色葡萄球菌 RN1304 及其变株的质粒消除与转导.中国药科大学学报,1989;20(3):151
- 4 陈小英,冯瑞山,陈知本.金黄色葡萄球菌抗生素、耐药质粒 DNA 的抽提及检测.南京药学院学报,1985;16(2):53.
- 5 Sambrook J, Fritsch E F, Maniatis T. Molecular Cloning 1989 Cold spring harbor laboratory. 6,30-31
- 7 王岳五,陈月华,陈宁,郭淑华,焦瑞身.球形芽孢杆菌 TS-1 与苏云金芽孢杆菌 H₄ 细胞融合的研究.生物工程学报,1988;4(3):230
- 8 Shing Chang and Conen S N. High frequency transformation of *Bacillus subtilis* Protoplasts. *Mol Gen Genet*, 1979;188:111
- 9 孙胜,袁丽蓉.生米卡链霉菌与北里链霉菌种间原生质体融合重组的研究.中国抗生素杂志,1989;14(5):344
- 10 Santamaria R, Gil J A and Martin J. High-frequency transformation of *Brevibacterium lactofermentum* protoplasts by plasmid DNA. *J Bacteriol*, 1985;162:463

Protoplast Transformation of Staphylococcal DNA

Feng Ruishan, Chen Xiaoying, Chen Zhiben

Biological Research Center

Experiment on protoplast-transformation of *S. aureus* RN1304(Y-1) by plasmid DNA or total DNA extracted from multiple antibiotic resistant staphylococcal strains (80022, 82022 and 82062) was reported. Protoplasts were prepared by treating recipients at early stationary phase with lysozyme and lysostaphin. Mixed protoplasts and DNA, then treated with polyethylene glycol (molecular weight. 6000), Tc^r or Cm^r transformants were obtained. The frequencies of protoplast transformation were 10^2 — 10^3 transformants per μg DNA.

Key words Staphylococcus; Protoplast-transformation; Plasmid DNA

【文摘044】复方新诺明片的生物利用度研究 邵俊,毛凤斐,屠锡德.中国医药工业杂志,1992;23(8):352

设计了速释片剂新处方 C,并建立了测定溶出介质和血浆中磺胺甲噁唑(SMZ)和甲氧苄啶(TMP)含量的反相高效液相色谱法。体外溶出度和人体内生物利用度的研究表明,C的主药吸收快,血浆峰浓度高,但血浓-时间曲线下面积与另两片剂 A 和 B 无显著差异。在人工胃液中,三种片剂 TMP 溶出均迅速,

但 SMZ 的平均溶出时间有显著差异。

【文摘045】对苯二胺的合成 莫芬珠,高林东,黄嘉梓.中国医药工业杂志,1992;23(8):370

以对苯二甲酸为起始原料,经酰化、氨化、降解三步反应合成对苯二胺。氨化可在较低压力和温度下完成;酯可用聚酯纤维生产中的废酯代替;降解反应将氮气保护下的密闭操作改为常规操作。三步总收率 69.5%,产品采用芳伯胺重氮法测定含量,达到国内质量标准。