

- tion of polydimethylsiloxane rubber in extracts of siliconized paper by IR spectrometry. *Zhong Lub*, 1987, 53(12), 25
- 7 张叔良,易大年,吴天明编著. 红外光谱分析与新技术. 北京:中国医药科技出版社,1993. 150
- 8 陈允魁编著. 红外吸收光谱法及其应用. 上海:上海交通大学出版社,1993. 97
- 9 董庆年编. 红外光谱法. 北京:石油化学工业出版社,1977. 188

## Determination of Polydimethylsiloxane in Meteospasmyl Capsules by IR

Zhang Yuying, Huang Jingyu, Hu Jiachi

Guangzhou Institute for Drug Control, Guangzhou 510160

**Abstract** This paper reports the determination of polydimethylsiloxane in Meteospasmyl capsules by IR. Polydimethylsiloxane was extracted from Meteospasmyl capsules with carbon tetrachloride. The solution was placed in a filled cell, the IR spectrum was recorded and the background was deducted with computer. Quantitative determination of polydimethylsiloxane was at  $1260 \text{ cm}^{-1}$ . A linear relation was obtained in the range of  $4 \sim 8 \text{ mg/ml}$ ,  $r = 0.9998$ . The mean recovery was 101.1% ( $n = 5$ ), and  $RSD = 2.0\%$ .

**Keywords** Meteospasmyl capsules; Polydimethylsiloxane; IR; Determination of content

**[文摘 013]** 正交实验设计筛选酒炙白芍的最佳炮制条件 杨中林,卢风兰,张琳等. 中药材,1996, 19(5):238

采用正交实验设计,以芍药甙的含量测定为指标,筛选酒炙白芍的炮制条件。其结果提示以加酒量5%,温度90℃,加热10 min为最佳炮制条件。

**[文摘 014]** 海龙科药用资源调查 张朝晖,徐国钧,徐培珊等. 中药材,1997, 20(5):224

对我国沿海海龙科药用动物资源调查结果表明,目前供药用的有7属12种,并就人工养殖问题作了分析。

**[文摘 015]** 吡乙酰胺合成工艺改进 王礼琛,芦金

荣,张德伟,黄嘉梓. 现代应用药学,1997, 14(1):29

以 $\alpha$ -吡咯烷酮和氯乙酸甲酯为原料,经缩合、氨解反应合成吡乙酰胺。并用正交设计和均匀设计试验优化反应条件,总收率达66%。

**[文摘 016]** 气质联用研究厚朴及其炮制品中挥发油 曾诠,周干南,屠万蔚等. 中草药,1997, 28(2):81

研究厚朴炮制前后挥发油的变化。结果表明:总含量降低,炒黄与姜炙厚朴相似,挥发油含量降低26%,炒焦降低42%,而其化学成分未发生明显变化,气相色谱图基本一致。实验发现挥发油中含有发汗、祛痰、平喘成分。